

The Core Whole Body Technique

Client: _____

Date: _____

Column II
IR Dosage DZ reading

Column I
IR Dosage DZ Reading

Column III
IR Dosage DZ Reading

Face 6 points

IR/Dosage IR/Dosage DZ reading

		**	_____
--	--	----	-------

CA		**	_____
----	--	----	-------

		**	_____
--	--	----	-------

		**	_____
--	--	----	-------

CB		**	_____
----	--	----	-------

		**	_____
--	--	----	-------

/	/	**	_____
---	---	----	-------

		**	_____
--	--	----	-------

Start at C7 first

A		**	_____
---	--	----	-------

		**	_____
--	--	----	-------

/	/	**	_____
---	---	----	-------

		**	_____
--	--	----	-------

B		**	_____
---	--	----	-------

		**	_____
--	--	----	-------

		**	_____
--	--	----	-------

C		**	_____
---	--	----	-------

		**	_____
--	--	----	-------

		**	_____
--	--	----	-------

D		**	_____
---	--	----	-------

		**	_____
--	--	----	-------

		**	_____
--	--	----	-------

E		**	_____
---	--	----	-------

		**	_____
--	--	----	-------

		**	_____
--	--	----	-------

F		**	_____
---	--	----	-------

		**	_____
--	--	----	-------

		**	_____
--	--	----	-------

G		**	_____
---	--	----	-------

		**	_____
--	--	----	-------

		**	_____
--	--	----	-------

H		**	_____
---	--	----	-------

		**	_____
--	--	----	-------

		**	_____
--	--	----	-------

I		**	_____
---	--	----	-------

		**	_____
--	--	----	-------

		**	_____
--	--	----	-------

J		**	_____
---	--	----	-------

		**	_____
--	--	----	-------

		**	_____
--	--	----	-------

K		**	_____
---	--	----	-------

		**	_____
--	--	----	-------

		**	_____
--	--	----	-------

L		**	_____
---	--	----	-------

		**	_____
--	--	----	-------

		**	_____
--	--	----	-------

M		**	_____
---	--	----	-------

		**	_____
--	--	----	-------

		**	_____
--	--	----	-------

N		**	_____
---	--	----	-------

		**	_____
--	--	----	-------

1. Mark the spine - length of the electrodes
 2. Take Initial Readings at spine and paraspinals
 3. Record IRs
 4. Treat to a D (Dose) the highest IR at each segment
 5. Do IRs on neck to C7 (usually two segments)
 6. Treat highest IRs to a D (Dose)
 8. Including neck and back pathways, compare doses on each pathway. Treat highest D (Dose) to a DZ 0 (Dose, Z, Zero)
 - ***If there is not a D on a pathway, do not DZ
 9. Stimulate for 2 minutes the highest DZ0 on back
- 6 points on Face:
1. Take and record IRs at positions 1 and 2
 2. Dose highest IR
 3. Do same for 3 and 4, and then 5 and 6
 5. Take highest dose on face to DZ0
 6. Compare DZ0 on back and face. If the face zero is higher than the back, set device to deep stimulate and treat the DZ0 for two minutes. If back is higher, you are complete.